

Pre-Surgical Hold Times

Antiplatelet Hold Times

Brilinta **5 days**

(ticagrelor)

Effient **7 days***

(prasugrel)

Plavix **5 days**

(clopidogrel)

NSAIDs**

Bayer, Ecotrin **7 days***

(aspirin)

Aleve

(naproxen)

Cambia or Zipsor

(diclofenac)

Daypro

(oxaprozin)

3 days

Motrin

(ibuprofen)

1 day

Day of surgery is day zero

Adapted from Lexicomp®, FDA Prescribing Information, UpToDate® and Drug.com Half Life/Clearance Calculator. **NOTICE:** This document is for general informational purposes only and is not intended and shall not be used as a substitute for professional medical advice, diagnosis, treatment and/or independent medical judgment. Optimum Independent Practice Association is not responsible for any reliance on any information in this document and/or any errors or omissions, results and/or consequences arising from or related to any adoption and/or use of such information.

Anticoagulant Hold Times

Bevyxxa **4 days**

(betrixaban)

Coumadin **highly variable** among patients
(warfarin) **bridge** if mechanical valve

Eliquis **2 days***

(apixaban)

Pradaxa **2 days** before if CrCl \geq 50ml/min
(dabigatran) **5 days*** before if CrCl < 50ml/min

Savaysa **2 days**

(edoxaban)

Xarelto **1 day** if normal renal function
(rivaroxaban) **2 days** if elderly or evidence of renal impairment

Day of surgery is day zero

*May hold longer if risk of bleed outweighs the benefit, in the providers professional opinion, based on the type of surgery (spinal surgery) and stroke risk.

**This is not an all-inclusive list

